

Eclampsia Checklist

- Call for Assistance
- Designate
 - Team leader
 - Checklist reader/recorder
 - Primary RN
- Ensure side rails up
- Protect airway and improve oxygenation:
 - Maternal pulse oximetry
 - Supplemental oxygen (100% non-rebreather)
 - Lateral decubitus position
 - Bag-mask ventilation available
 - Suction available
- Continuous fetal monitoring
- Place IV; Draw preeclampsia labs
- Ensure medications appropriate given patient history
- Administer magnesium sulfate
- Administer antihypertensive therapy if appropriate
- Develop delivery plan, if appropriate
- Debrief patient, family, and obstetric team

† "Active asthma" is defined as:

- (A) symptoms at least once a week, or
- (B) use of an inhaler, corticosteroids for asthma during the pregnancy, or
- (C) any history of intubation or hospitalization for asthma.

REVISED JULY 2017

MAGNESIUM SULFATE

Contraindications: Myasthenia gravis; avoid with pulmonary edema, use caution with renal failure

IV access:

- Load 4-6 grams 10% magnesium sulfate in 100 mL solution over 20 min
- Label magnesium sulfate; Connect to labeled infusion pump
- Magnesium sulfate maintenance 1-2 grams/hour

No IV access:

- 10 grams of 50% solution IM (5 g in each buttock)

ANTIHYPERTENSIVE MEDICATIONS

For SBP \geq 160 or DBP \geq 110

(See SMI algorithms for complete management when necessary to move to another agent after 2 doses.)

- Labetalol** (initial dose: 20mg); **Avoid parenteral labetalol with active asthma, heart disease, or congestive heart failure; use with caution with history of asthma**
- Hydralazine** (5-10 mg IV* over 2 min); **May increase risk of maternal hypotension**
- Oral Nifedipine** (10 mg capsules); Capsules should be administered orally, not punctured or otherwise administered sublingually

* Maximum cumulative IV-administered doses should not exceed 220 mg labetalol or 25 mg hydralazine in 24 hours

Note: If persistent seizures, consider anticonvulsant medications and additional workup

ANTICONVULSANT MEDICATIONS

For recurrent seizures or when magnesium sulfate contraindicated

- Lorazepam (Ativan):** 2-4 mg IV x 1, may repeat once after 10-15 min
- Diazepam (Valium):** 5-10 mg IV q 5-10 min to maximum dose 30 mg

FOR PERSISTENT SEIZURES

- Neuromuscular block and intubate
- Obtain radiographic imaging
- ICU admission
- Consider anticonvulsant medications