



The American College of
Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

ACOG
THE AMERICAN CONGRESS
OF OBSTETRICIANS
AND GYNECOLOGISTS

Statement of Policy

As Issued by the Executive Board of the
American College of Obstetricians and Gynecologists and the
American Congress of Obstetricians and Gynecologists

TOBACCO MARKETING AIMED AT WOMEN AND ADOLESCENTS

The American College of Obstetricians and Gynecologists and the American Congress of Obstetricians and Gynecologists find the targeting of women of all ages by the tobacco industry unconscionable and strongly oppose the practice.

The health risks of tobacco use disproportionately affect the health of women and are well documented (1,2,3). It also is well known that smoking by a pregnant woman is harmful to her fetus (4-12). Because of these well-known dangers, it is irresponsible for corporations involved in the production of tobacco products to single out women, especially those who are young, at risk for pregnancy, or educationally or otherwise disadvantaged, and encourage them to smoke.

Specifically, corporations involved in the production of tobacco and nicotine products (including, but not limited to cigarettes, pipes, cigars, cigarillos, little cigars, bidis, kreteks, tobacco [including chew, snuff], and powdered/dissolvable tobacco in the form of strips, sticks, or lozenges) and electronic nicotine delivery systems must stop targeting their marketing efforts, including advertising, market research, media planning, public relations, product pricing, distribution, customer support, sales strategy, and community involvement, to encourage women, particularly adolescents, to smoke cigarettes, or consume any form of tobacco product (13). The health of all women and future generations demands that consideration.

In accordance with our concerns about the negative effects of tobacco use, the American College of Obstetricians and Gynecologists and the American Congress of Obstetricians and Gynecologists will not accept funding for any purpose from corporations involved in the production of tobacco products.

References:

1. Office on Smoking and Health. Women and smoking: a report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention; 2001. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK44303/>. Retrieved July 20, 2016.
2. U.S. Department of Health and Human Services. How tobacco smoke causes disease: the biology and behavioral basis for smoking-attributable disease: a report of the Surgeon General. Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK53017/#!po=16.6667>. Retrieved July 20, 2016.

Tobacco Marketing Aimed at Women and Adolescents

Page 2

3. Tobacco use and women's health. Committee Opinion No. 503. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2011;118:746-50.
4. Tong VT, Dietz PM, Morrow B, D'Angelo DV, Farr SL, Rockhill KM, et al. Trends in smoking before, during, and after pregnancy--Pregnancy Risk Assessment Monitoring System, United States, 40 sites, 2000-2010. Centers for Disease Control and Prevention. *MMWR Surveill Summ* 2013;62:1-19.
5. Castles A, Adams EK, Melvin CL, Kelsch C, Boulton ML. Effects of smoking during pregnancy. Five meta-analyses. *Am J Prev Med* 1999;16:208-15.
6. Spinillo A, Nicola S, Piazzzi G, Ghazal K, Colonna L, Baltaro F. Epidemiological correlates of preterm premature rupture of membranes. *Int J Gynaecol Obstet* 1994;47:7-15.
7. Salihu HM, Aliyu MH, Pierre-Louis BJ, Alexander GR. Levels of excess infant deaths attributable to maternal smoking during pregnancy in the United States. *Matern Child Health J* 2003;7:219-27.
8. Sondergaard C, Henriksen TB, Obel C, Wisborg K. Smoking during pregnancy and infantile colic. *Pediatrics* 2001;108:342-6.
9. von Kries R, Toschke AM, Koletzko B, Slikker W, Jr. Maternal smoking during pregnancy and childhood obesity. *Am J Epidemiol* 2002;156:954-61.
10. England LJ, Kendrick JS, Wilson HG, Merritt RK, Gargiullo PM, Zahniser SC. Effects of smoking reduction during pregnancy on the birth weight of term infants. *Am J Epidemiol* 2001;154:694-701.
11. DiFranza JR, Aligne CA, Weitzman M. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics* 2004;113:1007-15.
12. Smoking cessation during pregnancy. Committee Opinion No. 471. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2010;116:1241-4.
13. U.S. Department of Health and Human Services. Preventing tobacco use among youth and young adults: a report of the surgeon general. Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2012. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK99237/>. Retrieved July 20, 2016.

Approved by the Executive Board July 1990
Reaffirmed July 2000
Revised and Approved July 2004
Reaffirmed July 2007
Revised and Approved July 2009
Revised and Reaffirmed July 2013
Revised and Reaffirmed July 2016